

**THE FIRST MEDITERRANEAN SYMPOSIUM
ON THE CORALLIGENOUS AND OTHER
CALCAREOUS BIO-CONCRETIONS**



**January 2009
Tabarka - Tunisia**

INDEX

I.	Deep sea coralligenous beds observed with ROV on four seamounts in the western Mediterranean	4
II.	Species distribution for Bio-concretions (maërl and coralligenous beds) and other ecosystems not associated with red algae	8
III.	Images from ROV dives on the seamounts in the study (Different stages of coralligenous concretions and species found on the seamounts researched)	9
IV.	Main threats and proposals for the protection of the coralligenous beds in the Mediterranean sea	13

I. DEEP-SEA CORALLIGENOUS BEDS OBSERVED WITH ROV ON FOUR SEAMOUNTS IN THE WESTERN MEDITERRANEAN

By Ricardo Aguilar, Xavier Pastor, Ana de la Torriente & Silvia García
OCEANA, Leganitos, 47. 28013 Madrid-SPAIN. raguilar@oceana.org

Abstract

*In 2006 and 2007, Oceana carried out several investigations on four Western Mediterranean seamounts, finding red algae bio-concretions down to 150-160 meters depth. The areas surveyed were the Ausias March seamount and the Emile Baudot seamount in the Mallorca Channel (Balearic Islands), the Chella Bank (Andalusia-Alboran Sea) and the Palos seamount (in front of Palos Cap, in Murcia). More than 40 hours of video material was collected with an ROV. Species known only to be in shallow waters, like carnivorous sponges (*Asbestopluma hypogea*), were found on small seamounts with peaks between 80 and 170 meters depth. Sponge aggregations were filmed on coralligenous beds and new data on the distribution of anthozoans (e.g., *Paramuricea macrospina*) was recorded. Nearly 300 species living on these bottoms were identified, giving new perspectives on their range and habitat dependence and preferences.*

Key words: seamounts, coralligenous, bio-concretions, maërl, carnivorous sponge.

Introduction

Red calcareous algae have been widely studied in the shallow waters of the Western Mediterranean (Ballesteros, 2006), but there is very little information about their distribution and function in deep areas. Two main infralittoral and circalittoral ecosystems created by calcareous red algae have been mentioned: maërl and coralligenous beds (Pérès & Picard, 1964; Picard, 1965). These have been described as areas of high diversity and ecological importance (Bosence, 1983; Barberá et al., 2003), being two of the most productive ecosystems in temperate regions (Martin et al., 2007). Seamounts and smaller marine elevations are considered hotspots, “stepping stones” and zones with high biodiversity (Matthiessen et al. 2003; Butler et al. 2001; Morato & Pauly (eds), 2004). Those with shallow peaks are often found to be areas of high biological productivity (Rogers 1994), as in the four seamounts researched, with tops between 80 and 100 m. depth, where red algae can grow and develop.

Materials and methods

The research was carried out during from June to September of 2006 and 2007 onboard the Oceana “Ranger” catamaran, equipped with a HSB2-plus Raymarine digital sonar with a high-powered transducer, linked to software to create bathymetric maps. Nineteen dives were carried out on four marine seamounts (Fig.1). Transects were filmed by a camera with 750 lines of resolution, a F1.2 lens and a 1:12 zoom, attached to an ROV Phantom H2+2. The ROV provided real time data on its position, depth, course, day and time. All of the identifications were made visually.

Results

Two main red algae formations were registered: (i) maërl or rhodolith beds and (ii) coralligenous formations. Most of the rhodolith beds found on these mounds and seamounts reached down to 140-150 meters depth, although the most important ones were between 80 and 120 meters. The formations were especially common over the top of Ausias March, but could also be found on Emile Baudot and the Chella Bank; they were absent from the Palos seamount. Three forms of coralligenous beds were detected: (i) large bio-concretions, (ii) “cobble” bio-concretions and (iii) thin sheets and small patches.

Although some smaller patches were found at 160-170 meters depth, large concretions were more common between 80 and 120 meters depth. Flat areas on the top of the seamounts showed the largest bio-concretions, normally formed by red calcareous algae of the genera *Lithophyllum*, *Mesophyllum* and *Neogoniolithon*, usually with other red algae, like *Peyssonnelia* spp. and the green algae *Palmophyllum crassum*. The most important ones were found on Ausias March and Chella Bank. Large bio-concretions forming round circles of around two meters in diameter and ten to 20 centimetres high were found on top of the Ausias March mound. These kinds of geometrical concretions were not found over the other seamounts. Coralligenous beds did not always form large bio-concretions but instead small, spotted blocks of some 10 to 30 centimetres in diameter fixed in the substratum. It was very often found as a transitory substratum between maërl and large coralligenous beds. They were very common on Ausias March and Emile Baudot. Patches of red algae were found on all of the seamounts. They were very common on the Palos mound, but were also the most common bio-concretion over the 120-130 meter range.

	Ausias March 38°44'N-001°48'E	Emile Baudot 38°42'N-002°20'E	Palos seamount 37°53'N-000°01'W	Chella Bank 36°31'N-002°51'W
Dives	3	4	6	6
Nautical miles	1.85	3.7	1,77	1.96
Area observed m ²	5,140	10,278	4,917	5,445
Filming time	5h29m	9h40m	8h24m	14h57m

Fig.1 Summary of dives, time and areas observed with the ROV on the four seamounts

Some 300 species were identified. 150 of them were most commonly found in red algae bio-concretions, but none of them were exclusive from these beds. Two biological communities were widely distributed on bio-concretion beds: sponge aggregation (genera *Haliclona*, *Aplysina*, *Tedania*, *Axinella*, etc.) and fields of dead man’s fingers (*Alcyonium palmatum* and *Paralcyonium spinulosum*). Species like *Paramuricea clavata*, *P. macrospina*, *Anthias anthias*, *Muraena helena*, *Lappanella fasciata* and *Phycis phycis* were recorded mainly on coralligenous beds. The carnivorous sponge *Asbestopluma hypogea* was first found in deep areas, but not always connected to bio-concretions. The specimen found on Ausias March was on a coralligenous bio-concretion at 100 meters depth, but the one found in Chella Bank was at 167 meters in a rocky area on a small pinnacle beside the main summit. Some other protected species included in the annexes of BAR-COM-SPAM were also found. For example, the elephant ear sponge (*Spongia agaricina*) was found on Emile Baudot and the triton snail (*Charonia lampas*) on Ausias March.

Discussion and conclusions

Maërl was mainly formed by rounded rodoliths, instead of the branched forms more common in shallower areas. Hydrodynamism and bathymetric distribution can determine morphology and maërl ramification (Bosence, 1983, Steller and Foster, 1995, Yabur-Pacheco and Riosmena-Rodríguez, 2007). Coralligenous concretions went from thin patches to large concretions -including circular geometric formations not yet described - many times looking like steps or visible different stages as it builds up. As Laborel (1961) affirms, morphology and interim structure could depend on depth, topography and algae species. Some concretions give an aspect of a cobbled seabed, likely due to the lack of fusion or coalescence between several patches of algae, as in the large bio-concretions. Although red algae bio-concretions were found in all of the areas researched, from the surface down to 160-170 meters depth, distribution of the communities had a spatial segregation.

Most of the species found associated with the coralligenous beds were also found in surrounding areas without red algae, including *Anthias anthias*, *Lappanella fasciata*, *Muraena helena* and *Phycis phycis*, although they were apparently less abundant, showing their preferences for irregular bottoms. Only a few species, like *Paramuricea clavata*, seem to be strongly related to these bio-concretions, although depth distribution is probably a more important factor. Sponge aggregations were more common on maërl and “cobbled” coralligenous beds, while dead man’s finger colonies were more often found on “cobbled” coralligenous, large coralligenous and rocky areas. *Asbestopluma hypogea*, since it was discovered in 1995 (Vacelet & Boury-Esnault, 1996), was so far only recorded in shallow caves in France and Croatia. Although Bakran-Petricioli et al. (2007), mentioned the possibility, this is the first time this species has been found in deep areas, both in coralligenous beds and rocky bottoms.

Acknowledgments

This work would not be possible without the financial help from Fundación Biodiversidad and La Obra Social de “La Caixa”.

Bibliography


- BALLESTEROS E. (2006) - Mediterranean Coralligenous assemblages: a synthesis of present knowledge. *Oceanogr. and Mar. Biol.: an Annual Review*. 44: 123-195.
- BAKRAN-PETRICIOLIT., VACELET J., ZIBROWIUS H., PETRICIOLI D., CHEVALDONNÉ P., RAĐA T. (2007). New data on the distribution of the 'deep-sea' sponges *Asbestopluma hypogea* and *Oopsacas minuta* in the Mediterranean Sea *Marine Ecology*, Volume 28, Supplement 1, September 2007, pp. 10-23(14).
- BARBERÁ C., BORDEHORE C., BORG J.A., GLÉMAREC M., GRALL J., HALL-SPENCER J.M., DE LA HOZ C.H., LANFRANCO E., LASTRA M., MOORE P.G., MORA J., PITA M.E., RAMOS-ESPLÁ A.A., RIZZO M., SÁNCHEZ-MATA A., SEVA A., SCHEMBRI P.J., VALLE C., (2003) Conservation and management of northeast Atlantic and Mediterranean maerl beds. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13: 65-76.
- BOSENCE D.W. (1983). Coralline algal reef framework. *J. Geol. Soc. London*: 140: 365- 376.
- BUTLER A.J., KOSLOW J.A., SNELGROVE P.V.R., JUNIPER S.K. (2001) A review of the biodiversity of the Deep Sea. Environment Australia, Canberra, 2001. Available from website: <http://www.ea.gov.au/marine>
- LABOREL J. (1961). Le concretionnement algal "coralligène" et son importance géomorphologique en Méditerranée. *Recueil Travaux Station Marine d'Endoume*, 23: 37-60.
- MARTIN S., CLAVIER J., CHAUVAUD L., THOUZEAU G. (2007). Community metabolism in temperate maerl beds. II. Nutrient fluxes. *Marine Ecology Progress Series* 335: 31-41.
- MATTHIESSEN B., FOCK H.O., VON WESTERNHAGEN H. (2003). Seamounts, hotspots for high speciation rates in benthic-pelagic fishes? A case study on *Macrorhamphosus* spp. (Syngnathidae) from the Great Meteor seamount, *Helgoland Marine Research*, 57, 63-72.
- MORATOT., PAULY D. (eds). 2004. Seamounts: Biodiversity and fisheries. *Fisheries Centre Research Reports* 12(5): 78pp.
- PÉRÈS J., PICARD J.M. (1964). Nouveau manuel de bionomie benthique de la mer Méditerranée. *Recueil Travaux Station Marine d'Endoume*, 31(47): 1-131.
- PICARD J. (1965). Recherches qualitatives sur les biocénoses marines des substrats meubles dragables de la région marseillaise. *Recueil des Travaux de la Station Marine d'Endoume*. 52(36): 1-160.
- STELLER D. L., FOSTER M. S. (1995). Environmental factors influencing distribution and morphology of rhodoliths in Bahía Concepción, B. C. S., México, *J. Exp. Mar. Biol. Ecol.* 194: 201-212.
- VACELET J., OURY-ESNAULT N. (1996). A new species of carnivorous sponge (Demospongiae: Cladorhizidae) from a Mediterranean cave. In: Willenz P, editor. Recent advances in sponge biodiversity inventory and documentation. *Bull Institut Roy Sci Natur Belg Sci Terre, Biologie*. 1996; 66 Suppl: 109-115.
- YABUR-PACHECO R. RIOSMENA-RODRÍGUEZ R. (2007). Rhodoliths beds composition in the southwestern Gulf of California, Mexico. *Publications of the Seto Marine Biological Laboratory Special Publication Series*. Selected Papers of the NaGISA World Congress 2006. Vol.8: 37-47 pp.

III. IMAGES FROM ROV DIVES ON THE SEAMOUNTS IN THE STUDY (DIFFERENT STAGES OF CORALLIGENOUS CONCRETIONS AND SPECIES FOUND ON THE SEAMOUNTS RESEARCHED)


Steps of different stages of coralligenous concretions found on the seamounts researched




maërl o rhodoliths bed



"cobbled" bio-concretions



rhodoliths-cobbled



thin sheets




transition from cobbled to large bio-concretion




large bio-concretion

Rhodoliths bed at 146 m of depth in Emile Baudot seamount.



Species recorded in deep areas of Ausias March seamount



anthozoans *Paramuricea macrospina*




carnivorous sponge
Asbestopluma hypogea

Protected species included in the annexes of BARCOM-SPAM recorded on the seamounts



triton snail (*Charonia lampas*)
in Ausias March



Elephant ear sponge (*Spongia agaricina*)
in Emile Baudot

Species widely distributed on bio-concretions bed found on the four seamounts researched



Dendrophyllia cornigera



Diazonia violacea



Palinurus elephas, *Paralcyonium spinulosum*
and unknown *nephteida*



Astroportus mediterraneus



Capros aper and *Pteroides griseum*




Phycis phycis



Callogorgia verticillata



Sponge not identified



Paramuricea clavata




Luidia ciliaris




Muraena helena in *Paramuricea clavata* garden




Aplysina aerophoba




Eunicella verrucosa




Paralcyonium spinulosum




Sponge field on maerl



Echinus melo



Calappa granulata



Viminella flagellum



Scorpaena sp.



Alcyonium palmatum

IV. MAIN THREATS AND PROPOSALS FOR THE PROTECTION OF CORALLIGENOUS BEDS IN THE MEDITERRANEAN SEA

During OCEANA's investigations large amounts of debris, lost or abandoned fishing gear and trawl marks were seen on the sea floor. Marine communities with high ecological value such as the red algae concretions are faced with numerous threats, especially when the constructing species have slow growth rates and are very long-lived. Some coralligenous concretions have been dated back to 8,000 years¹ and some rhodoliths can reach 100 years old².

It is known that different fishing techniques cause severe damage to the marine bottom. Both maërl and coralligenous beds are highly vulnerable to anthropogenic alterations. The use of trawls or dredges in fishing grounds can bring about irreparable damages, due to direct impacts on the coralligenous concretions and to resuspension of sediments that cause turbidity in the water and burial of the organisms. In some maërl communities destroyed by fishing, there are no recovery signs until some years later³.

In the Mediterranean Sea there are large amounts of debris, coming both from the earth and from recreational and commercial fishing boats. Fishing gear, plastics and other non-identified objects were found in many of the sampled areas.

Many boats anchor without any kind of control in all types of sea beds. In particular, frequent diving activities in areas of high ecological value can bring about serious damage from repeated anchoring in coralligenous zones.

It has been verified that a wide variety of coralligenous species is strongly affected by contamination, causing a loss of biological diversity⁴. The effect on constructing species is especially serious, as bad water quality can increase the number of organisms that erode red algae concretions⁵.

The introduction of exotic species into European ecosystems is one of the most worrisome threats for many marine communities. One of the invasive species that represents the greatest danger for coralligenous beds is *Womersleyella (Polysiphonia) setacea*, since it grows abundantly and reduces the amount of light that reaches the constructing algae⁶ and diminishes biodiversity in this community⁷. Other invasive species that can affect the growth and development of this habitat are *Lophocladia lallemandii*, *Asparagopsis taxiformis* and *Caulerpa taxifolia*.

Another serious threat for these ecosystems is climate change, as changes in sea temperatures can affect calcification rates of the algae. Similarly, diverse mortality episodes⁸ of suspension-feeding animals in hundreds of kilometres of coralligenous beds in the Mediterranean Sea has been related to high temperatures that were reached in these waters over long periods of time⁹.

Oceana proposes the following measures, which must be applied urgently, to protect coralligenous beds and end their current decline:

Regulation and control of the threats

Industrial and recreational fishing, the presence of debris, sport diving, uncontrolled anchoring by boats, deterioration in water quality by land, coastal and high seas-based activities, and an increasing abundance of invasive species are the main causes of coralligenous bed decline and loss in the Mediterranean Sea. These activities must be controlled to eliminate the severe impact they have on these ecosystems.

Mapping and investigation of the existing communities

Data on the location, extension and distribution of habitats are necessary to study the ecological and geomorphological processes that occur, as well as the degree of alteration that these habitats have suffered due to the impacts of human activities. The European Commission and Mediterranean countries acknowledge that there is a generalized lack of scientific data and suitable resources to realise the necessary scientific work.

Mediterranean countries must avoid the deterioration and alteration of habitats which can have an appreciable effect on certain species. Given the lack of habitat and species data, the process of selecting coastal and marine protected areas has been prevented.

A study of the state of these vulnerable habitats is necessary to elaborate effective solutions to stop their deterioration.

One of the main goals of mapping coralligenous bottoms is to identify their location and extension. This enables the application of both existing legislation and new measures for their conservation.

Application and improvement of effective legislation

Lithothamnion corallioides and *Phymatolithon calcareum* are included in Annex V of the EU Habitats Directive concerning species whose collection and exploitation in nature can be a management object. Red algae *Goniolithon byssoides*, *Lithophyllum lichenoide*, *Ptilophora mediterranea* and *Schimmelmannia shousboei* are included in Annex II of the Protocol concerning Specially Protected Areas¹⁰. Some cnidarians, such as *Corallium rubrum*, are also included in this Protocol as well as the Bern Convention¹¹, along with different sponges and other organisms typical of this community. Even so, these classifications are insufficient for the protection of these habitats when they are not effectively fulfilled and corresponding sanctions from infractions are not applied.

In the case of the EU, the lack of protection for certain sea beds is one of the most important deficiencies in the Habitats Directive (92/43/CEE). Its Annexes pose problems for the declaration of marine habitats. In them, coralligenous beds are not considered high-priority habitats, nor are any of their characteristic anthozoans. Although these formations are indirectly considered under the habitat definition of "1170 Reefs" in Annex I, revision and improvement of this and other definitions are necessary to guarantee effective protection of these marine communities.

In addition, coralligenous beds should not have to withstand the damage brought about by different fishing techniques such as the bottom trawling, since their use is prohibited on coralligenous and maërl bottoms according to the COUNCIL REGULATION (21 1967/2006 EC) of December 2006 concerning management measures of the sustainable exploitation of fishery resources in the Mediterranean Sea.

Declaration of Marine Protected Areas

Red algae concretions might be the most complex and diverse communities of the Mediterranean Sea. One way or another they are related to a great number of vulnerable habitats and represent great biological interest. The coralligenous bed is a collection of habitats strongly associated with gorgonian gardens, sponge fields and aggregations, caves and grottos, etc.

Various species that are characteristic of coralligenous formations, such as red and yellow gorgonians, suffer serious damage due to frequent underwater dives in certain areas¹².

Some species that live in association with coralligenous beds are considered threatened or endangered in the Mediterranean Sea¹³, including the algae *Chondrymenia lobata*, *Halarachnion ligulatum*, *Halymenia trigona*, *Platoma cyclocolpa*, *Nemastoma dichotomum*, *Ptilophora mediterranean*, *Schizymenia dubyi* and *Laminaria rodriguezii*, as well as some molluscs like the date mussel (*Lithophaga lithophaga*), echinoderms like the long-spined sea urchin (*Centrostephanus longispinus*), and fish such as *Sciaena umbra* and *Umbrina cirrhosa*¹⁴.

Therefore, there is an urgent necessity to establish measures through the designation of new protected areas for benthic communities of interest that are threatened, like coralligenous and maërl bottoms, among others. By creating Marine Protected Areas wherever coralligenous formations occur, they would protect directly these and their associated habitats, and indirectly the large numbers of species that depend on them.

Bibliography

- 1 SARTORETTO S., VERLAQUE M., LABOREL J. (1996). Age of settlement and accumulation rate of submarine "Coralligène" (-10 to -60 m) of North Western Mediterranean Sea; relation to Holocene rise in sealevel. *Marine Geology*, 130 : 317-331.; SARTORETTO S. (1994). Structure et dynamique d'un nouveau type de bioconstruction à *Mesophyllum lichenoides* (Ellis) Lemoine (Corallinales, Rhodophyta). *Comptes Rendus Académie des Sciences Série III, Life Sciences* 317: 156-160.
- 2 ADEY W. H., MCKIBBIN D. L. (1970). Studies on the maërl species *Phymatolithon calcareum* (Pallas) nov. comb. and *Lithothamnium corallioides* Crouan in the Ria de Vigo. *Bot. Mar.*, 13: 100-106; FAZAKERLEY H., GUIRY M. D. (1998). The distribution of maerl beds around Ireland and their potential for sustainable extraction: phycology section. Report to the Marine Institute, Dublin. National University of Ireland, Galway.
- 3 FOSTER M.S. (2001). Rhodoliths: between rocks and soft places. *Journal of Phycology* 37:1-9; BIRKETT D. A., MAGGS C. A., DRING M. J. (1998). Maërl (Volume V). An overview of dynamic and sensitivity characteristics for conservation management of marine SACs. Scottish Association for Marine Science. (UK Marine SACs Project). 116 pages. PANAYOTIDIS, P., SEMROUD, R., SINNASSAMY, J.M., SPAN A., VUIGNIER G. (1991) - Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée : IUCN et RAC/SPA
- 4 PIAZZI L., PARDI G., BALATA D., CECCHI E., CINELLI F. (2002). Seasonal dynamics of a subtidal north-western Mediterranean macroalgal community in relation to depth and substrate inclination. *Botanica Marina*, 45: 243-252.
- 5 HONG J.S. (1983). Impact of the pollution on the benthic community. Environmental impact of the pollution on the benthic coralligenous community in the Gulf of Fos, northwestern Mediterranean. *Bulletin Korean Fisheries Society*, 16(3): 273-290.

- 6 PEREZT., GARRABOU J., SARTORETTO S., HAMELIN J.G., FRANCOUR P., VACELET J. (2000). Mortalité massive d'invertébrés marins: un événement sans précédent en Méditerranée nordoccidentale. *Comptes Rendus Académie des Sciences Série III, Life Sciences*, 323: 853-865.
- 7 AIROLDI L., RINDI F., CINELLI F. (1995). Structure, seasonal dynamics and reproductive phenology of a filamentous turf assemblage on a sediment influenced, rocky subtidal shore. *Botanica Marina*, 38: 227-237.
- 8 BALLESTEROS, E. (2003). The coralligenous in the Mediterranean Sea. Definition of the coralligenous assemblage in the Mediterranean, its main builders, its richness and key role in benthic ecology as well as its threats. Project for the preparation of a Strategic Action Plan for the Conservation of the Biodiversity in the Mediterranean Region (SAP BIO). RAC/SPA- Regional Activity Centre for Specially Protected Areas.; MAYOL J., GRAU A., RIERA F., OLIVER, J. (2000). Llista vermella dels peixos de les Balears. *Quaderns de Pesca*, 4: 1-126.
- 9 VACELET J. (1991). Report of a mission in Tunisia, Syria, Cyprus, Greece and Turkey in the context of the programme "Fight against the epidemic decimating sponges in the Mediterranean" Rome: FAO; Technical cooperation programme; PEREZT., GARRABOU J., SARTORETTO S., HAMELIN J.G., FRANCOUR P., VACELET J. (2000). Mortalité massive d'invertébrés marins: un événement sans précédent en Méditerranée nordoccidentale. *Comptes Rendus Académie des Sciences Série III, Life Sciences*, 323: 853-865.
- 10 COMA R., POLÀ E., ZABALA M., ROS J. (2001). Effect of recreational activity on the coralligenous community: a decade of assessment of demographic parameters of the gorgonian *Paramuricea clavata*. Poster presentation. European Marine Biology Symposium. Maó, Menorca.
- 11 Convention on the conservation of European wildlife and natural habitats (Bern Convention). DO L 38 de 10.2.1982, p. 3/32
- 12 BOUDOURESQUE C. F., BALLESTEROS E., BEN MAIZ N., BOISSET F., BOULAUDIER E., CINELLI F., CIRIK S., CORMACI M., JEUDY DE GRISSAC A., LABOREL J., LANFRANCO E., LUNDBERG B., MAYHOUB H., MEINESZ A., PANAYOTIDIS P., SEMROUD R., SINNASSAMY J. M., SPAN A., VUIGNIER G. (1991) - Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée : IUCN et RAC/SPA Salammbô, G.I.S.-Posidonie, Marseille, Fr. : 1-250.
- 13 Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée PNUE. Plan d'Action pour la Méditerranée. Aires Spécialement Protégées (SPA), Alliance Mondiale pour la Nature (UICN), Groupement d'Intérêt Scientifique Posidonie, Athens 1990, UNEP/MAP
- 14 BARCOM (1995). The Protocol Concerning Mediterranean Specially Protected Areas (SPA Protocol); adopted in Geneva Switzerland, on 2 April 1982, in force 1986, revised in Barcelona, Spain on 9-10 June 1995 as the Protocol Concerning Specially Protected Areas and Biological Diversity in the Mediterranean (SPA and Biodiversity Protocol); EC (1999). 1999/800/EC: Council Decision of 22 October 1999 on concluding the Protocol concerning specially protected areas and biological diversity in the Mediterranean, and on accepting the annexes to that Protocol (Barcelona Convention).
- 15 BOUDOURESQUE C. F., BALLESTEROS E., BEN MAIZ N., BOISSET F., BOULAUDIER E., CINELLI F., CIRIK S., CORMACI M., JEUDY DE GRISSAC A., LABOREL J., LANFRANCO E., LUNDBERG B., MAYHOUB H., MEINESZ A., PANAYOTIDIS P., SEMROUD R., SINNASSAMY J.M., SPAN A., VUIGNIER G., (1991) - Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée : IUCN et RAC/SPA Salammbô, G.I.S.-Posidonie, Marseille, Fr. : 1-250. Boudouresque C.F., Avon M. & V. Gravez (eds.) 1991. Les espèces marines à protéger en Méditerranée. GIS Posidonie, Marseille. 447 pp.

PLaza de España - Leganitos, 47
28013 Madrid (Spain)
Tel.: +34 911 440 880
Fax: +34 911 440 890
e-mail: europa@oceana.org
www.oceana.org

Rue Montoyer, 39
1000 Brussels (Belgium)
Tel.: +32 (0) 2 513 22 42
Fax: +32 (0) 2 513 22 46
e-mail: europa@oceana.org

1350 Connecticut Ave. NW, 5th Floor
Washington, D.C. 20036 USA
Tel.: +1 (202) 833-3900
Fax: +1 (202) 833-2070
e-mail: info@oceana.org

175 South Franklin Street
Suite 418
Juneau, Alaska 99801 USA
Tel.: +1 (907) 586-4050
Fax: +1 (907) 586-4944
e-mail: northpacific@oceana.org

Avenida General Bustamante 24 Oficina 2C
Providencia, Santiago, Chile
Tel.: +56-2-795-7140
Fax.: +56-2-795-7146

